Journal of Immunological Methods, 163 (1993) 279–280
Elsevier Science Publishers B.V.

Contents to volume 163 (1993)

Number 1 – 6 July 1993

Immunological standardization

Nomenclature for human complement component C2. Who-IUIS Nomenclature Sub-Committee 1

Revised nomenclature for human complement component C4. WHO-IUIS Nomenclature Sub-Committee 3

Nomenclature for human complement factor B. WHO-IUIS nomenclature sub-committee 9

Research reports

Application of an immortalized human endothelial cell line to the leucocyte : endothelial adherence assay
K.A. Brown, A. Vora, J. Biggerstaff, C.-J.S. Edgell, S. Oikle, G. Mazure, N. Taub, A. Meager, T. Hill, C. Watson and D.C. Dumonde (UK, USA) 13

In vivo detection of fluorescent tumor-specific cytotoxic T cell clones
R.E.M. Toes, J.M.M. Den Haan, M.C.W. Feltkamp, R.J.J. Blom, C.J.M. Melief, E. Claassen and W.M. Kast (Netherlands) 23

Immortalizaton of antigen selected B cells
P.G.A. Steenbakkers, P.M.G.F. Van Wezenbeek and W. Olijve (Netherlands) 33

Competitive enzyme immunoassay for quantification of the cellular form of fibronectin (EDAcFN) in blood samples
S. Ylätupa, P. Partanen, C. Haglund and I. Virtanen (Finland) 41

A rapid biochemical method for measuring antigen-induced pulmonary eosinophil margination in allergic guinea pigs
P. Tagari, C. Black, S. Marshall and A.W. Ford-Hutchinson (Canada) 49

A sensitive enzyme immunoassay for the measurement of low concentrations of IgA
M. Hirvonen, S. Koskinen and H. Tölö (Finland) 59

Application of a monoclonal antibody against a neoepitope on activated C4 in an ELISA for the quantification of complement activation via the classical pathway
G.J. Wolbink, J. Bollen, J.W. Baars, R.J.M. Ten Berge, A.J.G. Swaak, J. Paardekooper and C.E. Hack (Netherlands) 67

Production and characterization of monoclonal antibodies raised against recombinant human granzymes A and B and showing cross reactions with the natural proteins
J.A. Kummer, A.M. Kamp, M. Van Katwijk, J.P.J. Brakenhoff, K. Radošević, A.M. Van Leeuwen, J. Borst, C.L. Verweij and C.E. Hack (Netherlands) 77

Enzyme-linked immunosorbent assay for quantification of anti-lipid A antibody. Enhancement of sensitivity by antigen immobilization at low pH
S.A. David and V.I. Mathan (India) 85

Hybrid hybridomas producing bispecific antibodies to CEA and peroxidase isolated by a combination of HAT medium selection and fluorescence activated cell sorting
P. Jantscheff, L. Winkler, L. Karawajew, G. Kaiser, V. Böttger and B. Micheel (Germany, UK) 91

Methods for computer assisted analysis of lymphoid cell shape and motility, including Fourier analysis of cell outlines
H. Verschueren, B. Houben, J. De Braekeleer, J. De Wit, D. Roggen and P. De Baetselier (Belgium) 99

Effect of red cell lysis protocols on the expression of rat peripheral blood lymphocyte subset and activation antigens
G.A. Webster and A.G. Pockley (UK) 115

A novel approach to the laboratory diagnosis of *Chlamydia trachomatis* infections using monoclonal anti-idiotypic antibodies
C. Laferrière, R.W. Peeling, E.S. Tackaberry, J. Hamel, J.-A. Dillon and B.R. Brodeur (Canada) 123

Letters to the editors

Isolation of rat blood lymphocytes using a two-step Percoll density gradient. Effect of activin (erythroid differentiation factor) on peripheral T lymphocyte proliferation in vitro
M.P. Hedger and L. Clarke (Australia) 133

Some comments on the Pick and Mizel microassay method for measurement of hydrogen peroxide production by macrophages
E. Hoffer, A. Tabak, L. Shenker and N. Lanir (Israel) 137

Corrigendum
In vitro phagocytosis inhibition assay for the screening of potential candidate antigens for sub-unit vaccines against the asexual blood stage of *Plasmodium falciparum*. (Journal of Immunological Methods, 159 (1993) 209–219)
J. Gysin, S. Gavoille, D. Mattei, A. Scherf, S. Bonnefoy, O. Mercereau-Puijalon, T. Feldmann, J. Kun, B. Müller and L.P. Da Silva (French Guiana, France, Germany) 139

Announcement 141

Notes to authors 143

Number 2 – 9 August 1993

Research reports
The release of soluble p55 TNF receptor from U937 cells studied by a new p55 immunoassay
N.-B. Liabakk, A. Sundan, E. Lien, A. Lægreid, M.P. Bombara, G. Ranges and T. Espevik (Norway, USA) 145
A simple flow cytometric procedure for the determination of surface antigens on unfixed leucocytes in whole blood
D.A. McCarthy and M.G. Macey (UK) 155
Enhancement of immunoassay sensitivity by molecular modification of competitors
R. Jockers, F.F. Bier and R.D. Schmid (Germany) 161
Two-site ELISA for quantification of the terminal C5b-9 complement complex in plasma. Use of monoclonal and polyclonal antibodies against a neoantigen of the complex
A. Accardo-Palumbo, G. Triolo, D. Casiglia, L. Salfi, E. Giardina and G. Triolo (Italy) 169
Triple immunofluorescence flow cytometry, using whole blood, of $CD4^+$ and $CD8^+$ lymphocytes expressing CD45RO and CD45RA
R. Gruber, C. Reiter and G. Riethmüller (Germany) 173
A fluorometric assay for the quantitation of cell adherence to endothelial cells
A.L. Akeson and C.W. Woods (USA) 181
Immunoassay reagents for psychoactive drugs. II. The method for the development of antibodies specific to imipramine and desipramine
M. Adamczyk, J. Fishpaugh, C. Harrington, D. Johnson and A. Vanderbilt (USA) 187
Human IgG in immune complexes bound to human erythrocyte CR1 is recognized differently than human IgG bound to an erythrocyte surface antigen
C.J. Reist, J.D. Wright, R.H. Labuguen and R.P. Taylor (USA) 199
A europium fluoroimmunoassay for measuring binding of antigen to class II MHC glycoproteins
S.M. Tompkins, P.A. Rota, J.C. Moore and P.E. Jensen (USA) 209
Preparation of monoclonal antibodies against the IL-6 signal transducer, gp130, that can inhibit IL-6-mediated functions
T. Saito, T. Taga, D. Miki, K. Futatsugi, H. Yawata, T. Kishimoto and K. Yasukawa (Japan) 217
The optimisation of a novel iodide microassay and its application in an immunoassay for human antibody levels in serum
R. O'Kennedy and P. Keating (Ireland) 225
Production of monoclonal and polyclonal antibodies against human osteocalcin sequences and development of a two-site ELISA for intact human osteocalcin
I. Kuronen, H. Kokko and M. Parviainen (Finland) 233
Use of density perturbation to isolate immunologically distinct populations of cells
D. Patel, C.P. Rubbi and D. Rickwood (UK) 241
A method for the detection of erythrocyte-bound interleukin-8 in humans during interleukin-1 immunotherapy
H. Tilg, D. Pape, E. Trehu, L. Shapiro, M.B. Atkins, C.A. Dinarello and J.W. Mier (USA) 253
Novel method for the measurement of cytokine production by a one-stage procedure
D. De Groote, Y. Gevaert, M. Lopez, R. Gathy, F. Fauchet, I. Dehart, M. Jadoul, D. Radoux and P. Franchimont (Belgium) 259

Short communication
Detection of the C3a complement component in commercial gamma globulins by dot blotting
R. Nezlin (Israel) 269

Letters to the editors
Evaluation of the methodology of polymorphonuclear leukocyte chemotaxis
J.L. Haddox and R.R. Pfister (USA) 273
Background noise in ELISA procedures. Influence of the pH of the coating buffer and correlations with serum IgM concentration
L.S. Kuen, C.H. Ming and Y.S. Fan (Malaysia) 277

Contents to volume 163 (1993) 279

Notes to authors 281